

Chinook Salmon Monitoring in the Lower Cowlitz



John Serl, Erick Rockwood, and Kale Bentley
Cowlitz Annual Program Review and Science Conference




Washington
Department of
**FISH and
WILDLIFE**

Washington Department of Fish & Wildlife
August 22nd, 2023

Photo credit: E Rockwood

Background

- Estimates of fall Chinook abundance in the lower Cowlitz R. have been generated since the 1960s
- Since 2010, Chinook have been monitored with aerial redd counts and carcass surveys to estimate abundance & composition
- In 2021, mark-recapture (M-R) carcass surveys were implemented in the lower river



Objectives

- Conduct carcass surveys w/ M-R
 - Obtain accurate abundance estimates
 - Estimate precisions
- Conduct aerial flights
 - Continue existing time series of abundance
 - Data for future bias-correction



Methods

- Aerial flights for redds
 - Timing: bi-weekly (scheduled late Sept. – early Dec.)
 - Space: Castle Rock to Barrier Dam (~33 miles)
 - Approach: Count & GPS all redds via helicopter
- Carcass surveys
 - Timing: Weekly (Sept. – Dec.); 4 – 5 days/week
 - Space: Olequa Ck to Barrier Dam (~26 miles)
 - Approach:
 - Jet boat + gaffes + CWT wand
 - 3 people & 1 boat
 - Recover all carcasses
 - Sample & tag representatively



Results: Aerial flights for redds

- Total Redds by Date

Date	Redds
14-Sep	39
28-Sep	393
12-Oct	849
26-Oct	1,152
9-Nov	-
22-Nov	-

- Abundance


- Spring-run

- Spawners: **111** (39 redds x 2.84 fish/redd)
- pHOS = 50% (59/119 HOR carcasses)

- Fall-run

- Spawners: **3,272** (1,152 redds x 2.84 fish/redd)
- pHOS = 10% (271/2,848 HOR carcasses)


- Total Spawners: **3,383**



Results: M-R surveys

- Surveys
 - 50 days across 18 weeks (Sept. 1st – Jan. 5th)
 - No missed survey weeks!
- Carcasses
 - Maiden (unique) = 3,015
 - Tagged = 1,178
 - Recaptured = 276

} Overall recovery probability ~30%
- Abundance
 - Total = **10,450** (median: 95% CI 8,791 – 13,480)
 - Spring-run
 - Spawners: **284** (median: 95% CI 184 – 493)
 - pHOS: 46%
 - Fall-run
 - Spawners: **10,150** (median: 95% CI 8,499 – 13,185)
 - pHOS: 7%




Results: 2021 vs. 2022


	Metric	2021	2022	Difference
Obj. #1	Carcasses maidens	2,167	3,015	+40% (848)
	Carcasses tagged	903	1,178	+30% (275)
	Carcasses recaptured	189	276	+46% (87)
	Recovery probability	~28%	~30%	+2%
	Fall Abundance: M-R	7,827	10,150	+30% (2,323)
Obj. #2	Fall Abundance: redds	4,913	3,272	-33% (-1,641)
	Ratio (redds/M-R)	0.63	0.32	-49% (-0.31)

Results: Redd Counts by Year


2021 & 2022



All Years



Years w/ "tail"



Conclusions

- Meeting objectives of the project
 - Obtain accurate estimates via M-R carcass surveys
 - Maintaining redd-based estimates for bias-correction later
- Next steps:
 - Short term → continue concurrent surveys
 - Long term → carcass tagging or updated expansion



Acknowledgments

- Survey crew
 - Erick Rockwood
 - Mike Blankenship
 - Nels Parvi
 - Carson Swart
- Helicopter flights
 - Northwest Helicopters
 - Chris Gleizes
- Funding
 - Tacoma Power
- Study Design & Analysis
 - Thomas Buehrens
 - Dan Rawding
- Data Management
 - Danny Warren






Questions?

Supplemental slides

Mark-Recapture: Data Collection & Analysis

Carcass Survey Flow Diagram



Abundance and composition of adult Chinook escapement is estimated using an **“open” population Jolly-Seber (JS) model** (Seber 1982, Pollock et al. 1990).

- “super population” JS model was developed by Schwarz et al. (1993, 1996) specifically for estimating salmon spawning escapement using mark-capture methods
- Has been successfully implemented to estimate spawner escapement for other salmon populations within the Lower Columbia River (Rawding et al. 2014) and other Washington state watersheds (Ashcraft et al. 2017).


Assumptions of Jolly-Seber Mark-Recapture

- ***Spatial and temporal coverage***: Carcasses are sampled and marked throughout the entire spawning run and encompass the entire spawning distribution.
- ***Equal Catchability***: Each carcass that is present in the study system during a specific sample event, whether tagged or untagged, has the same probability of being sampled.
- ***Equal Persistence***: Each carcass that is present in the study system during a specific sample event, whether tagged or untagged, has the same probability of survival (i.e., persisting in the study area to the following sample period).
- ***Tag Loss and Recovery***: Tagged carcasses do not lose their tags and all tags are recognized and read properly on recovery.
- ***Instantaneous Sampling***: All samples are instantaneous, i.e., the sampling time is negligible and each release is made immediately after the sample.

Flows in the lower Cowlitz for 4 years of redd-counts with “descending limb”

2018
doesn't
match the
pattern of
high flows
in early
Nov. that
could
have
resulted
in an
artificially
earlier
timed
peak
count

Earlier observed peak in redds



Later observed peak in redds

